

Tethered Interleukin-15 (IL-15)/IL-21 to Enhance T Cells for Cellular Therapy

Summary (1024-character limit)

Researchers at the National Cancer Institute (NCI) have developed a method to improve the function of therapeutic engineered T cells used for Adoptive T Cell Therapy (ACT) for various cancers and diseases through the co-expression of Interleukin-15 (IL-15) and IL-21 by a flexible linker to the cell membrane. Researchers at the NCI seek licensing and/or co-development research collaborations for this invention.

NIH Reference Number

E-068-2018

Product Type

Therapeutics

Keywords

 Interleukin-15, II-15, Interleukin-21, IL-21, T Cells, Adoptive Cell Therapy, ACT, Chimeric Antigen Receptor, CAR, Cancer, Hinrichs

Collaboration Opportunity

This invention is available for licensing and co-development.

Contact

John D. Hewes
 NCI - National Cancer Institute

240-276-5515

John.Hewes@nih.gov

Description of Technology

Interleukin-15 (IL-15) and IL-21 have been reported to support the function of anti-tumor T cells. However, their use in the clinic has been constrained, in part, by dose-limiting toxicity and the need for repeated administration. To overcome these limitations, researchers in the National Cancer Institute (NCI) Experimental Transplantation and Immunology Branch (ETIB) have developed synthetic IL-15 and IL-21 molecules for autocrine expression by the engineered therapeutic T cells. These molecules were designed with flexible linkers that connect to cell membrane anchors. This, in turn, reduces systemic toxicity caused by free cytokine molecules. The inventors have shown that co-expression of the novel IL-15 and IL-21 tethered molecules improves the anti-tumor efficacy of the therapeutic engineered T cells in vivo.



Potential Commercial Applications

• Treatment of cancer patients receiving T cell-based immunotherapy

Competitive Advantages

- T cells that co-express the tethered IL-15 and IL-21 on their cell membrane can increase therapeutic
 effectiveness of adoptive immunotherapy because it can reduce systemic toxicity caused by free
 cytokine molecules
- T cells that co-express the tethered IL-15 and IL-21 on their cell membrane are already known to have a greater decrease in tumor size compared to those mice treated with T cell-based immunotherapies using unmodified T cells

Inventor(s)

Christian Hinrichs (NCI), Benjamin Jin (NCI)

Development Stage

• Pre-clinical (in vivo)

Patent Status

• U.S. Provisional: U.S. Provisional Patent Application Number 62/628,454, Filed 09 Feb 2018

Therapeutic Area

Cancer/Neoplasm