

Wu et al., Mutational Analysis of the Adeno-Associated Virus Type 2 (AAV2) Capsid Gene and Construction of AAV2 Vectors with Altered Tropism, 2000, Journal of Virology, vol. 74, No. 18, pp. 8635-8647. cited by examiner .
GenBank Accession # AF513852, Adeno-associated virus 8 nonstructural protein and capsid protein, 2002. cited by examiner .
GenBank Accession # AY530579, Adeno-associated virus 9 isolate hu.14 capsid protein VP1 (cap), 2004. cited by examiner .
GenBank Accession # AF043303, Adeno-associated virus 2, complete genome, 1998. cited by examiner .
Appendix 1--Sequence alignment conducted on Apr. 2, 2008. cited by other .
Appendix 2--Sequence alignment conducted on Apr. 2, 2008. cited by other .
Bowles et al., "Marker rescue of adeno-associated virus (AAV) capsid mutants: a novel approach for chimeric AAV production", Journal of Virology, 77(1):423-432 (2003). cited by other .
Chen et al., "Molecular characterization of adeno-associated viruses infecting children", Journal of Virology, 79(23):14781-14792 (2005). cited by other .
Choi et al., "AAV hybrid serotypes: improved vectors for gene delivery", Current Gene Therapy, 5(3):299-310 (2005). cited by other .
Database UniProt [Online] "Capsid Protein VP1", retrieved from EBI Accession No. Uniprot:Q670R0, Database Accession No. Q670R0 (Oct. 11, 2004). cited by other .
Gao et al., Genbank Accession# AAS99284.1, Jun. 24, 2004. cited by other .
Gao et al., "Glades of Adeno-Associated Viruses Are Widely Disseminated in Human Tissues", Journal of Virology, 78(12): 6381-6388 (2004). cited by other .
Grimm et al., Journal of Virology, 80(1):426-439 (2006). cited by other .
International Search Report and Written Opinion for PCT/US2007/008314, Search report dated May 16, 2008, 17 pages. cited by other .
Ruffing et al., "Mutations in the carboxy terminus of adeno-associated virus 2 capsid proteins affect viral infectivity: lack of an RGD integrin-binding motif", Journal of General Virology, 75:3385-3392 (1994). cited by other .
Ruffing et al., Genbank Accession# AAC03780, Feb. 24, 1998. cited by other.

Primary Examiner: Lucas; Zachariah

Assistant Examiner: Blumel; Benjamin P

Attorney, Agent or Firm: Mohr; Judy M. Myers Fitch; Susan J. King & Spalding LLP

Government Interests

STATEMENT REGARDING GOVERNMENT INTEREST

This work was supported in part by the National Institutes of Health (NIH) Grant numbers HL 064274 and HL 066948. Accordingly, the United states government has certain rights.

Parent Case Text

CROSS-REFERENCE TO RELATED APPLICATIONS

the AAV-2 prototype at any dose and expressed over 100% of normal hFIX levels from intravenous injection of 5.times.10.sup.10 particles, whereas AAV-2 expression was over 100% of normal hFIX levels only at a dose of 1.times.10.sup.12.

In another study, recombinant human alpha-1-antitrypsin (hAAT)-expressing AAVs were prepared, from the AAV-DJ, AAV-2, AAV-8, or AAV-9 capsids. The hAAT gene was under an RSV promoter. Mice (C57/BL6) were injected via tail vein infusions of 2.times.10.sup.11 particles and plasma levels of hAAT were determined via specific ELISA 3, 7, and 14 days after injection. Results are shown in FIG. 5. AAV-8, AAV-9, and AAV-DJ expressed efficiently and equally outperformed the vector with an AAV-2 capsid.

In another in vivo study, liver transduction in the presence of human serum was quantified, to assess the ability of AAV-DJ to evade neutralization in vivo. As described in Example 4, mice were passively immunized with 4 or 20 mg IVIG prior to infusion of hFIX-expressing AAV-2, -8, -9, or -DJ. Plasma hFIX levels for each AAV serotype are shown in FIGS. 6A-6B as percent corresponding virus level in control mice treated with phosphate-buffered saline rather than IVIG as a function of time post infusion. FIG. 6A shows the results for mice immunized with 4 mg IVIG and FIG. 6B shows the results for mice immunized with 20 mg IVIG. AAV-2 expression was completely abolished, however transduction with AAV-DJ, -8 or -9 was inhibited in a dose-dependent manner, with AAV-DJ showing intermediate resistance at the high, and efficient evasion (similar to AAV-8 and AAV-9) at the low IVIG dose (FIG. 6A). These results were confirmed with a second independent IVIG batch from another vendor (Carimune 12%, Behring AG, data not shown).

In another study, also described in Example 4, the feasibility to repeatedly administer the different viruses to mice was assessed, to evaluate capsid cross-neutralization. Results are shown in FIG. 6C. No gene expression upon re-infusion of any of the capsids into animals already treated with the same serotype was observed. However, AAV-8 and -9 also efficiently blocked each other, substantiating previous data (Gao, G. et al., *J. Virol.*, 78:6381-6388 (2004)). This result might argue against the use of vectors based on these wildtypes in re-administration protocols, albeit they could be combined with AAV-2. In contrast, primary infusion of AAV-DJ allowed subsequent expression (up to 18%) from AAV-2, -8 or -9, likely due to the fact that AAV-DJ only shares a limited number of epitopes with each wildtype virus. In the reverse experiment, AAV-DJ vectors were inhibited in animals immunized with AAV-8 or -9, while giving detectable expression in AAV-2-treated mice. This implied a stronger or broader immune response from primary infusion of serotypes 8 or 9. AAV-DJ was more resistant to the corresponding mouse sera in culture, as seen in FIG. 6D. Less cross-reactivity between AAV-8 and -9 was noted.

AAV-DJ, as well as other recombinant protein capsids identified in the library discussed below, retained a heparin binding domain (HBD) from the AAV-2 parent. This domain functions in binding to the primary AAV-2 receptor heparin sulfate proteoglycan (Summerford, C. et al, *J. Virol.*, 72:1438-1445 (1998)). To investigate the role of the AAV-DJ HBD, two crucial arginine residues (Kern, A. et al., *J. Virol.*, 77:11072-11081 (2003)) were mutated to the respective residues in AAV-8 or -9, as shown in FIG. 7A, and are referred to herein as AAV-DJ/8 and AAV-DJ/9. Table 1 above includes data on the mutant AAV-DJ/8, and shows that gfp expression was reduced by several orders of magnitude, and was as low as that observed with serotypes AAV-8 or AAV-9.

DNA Family shuffling was used to create a complex library of hybrid particles from eight different wildtypes. Serial amplification on human cells enriched hybrids from a multitude of AAV serotypes, typically containing an AAV-2 heparin binding domain (HBD). More stringent selection with pooled human antisera yielded a single AAV-2-8-9 chimera, referred to herein as AAV-DJ. Recombinant AAV-DJ vectors were superior to natural AAVs in cultured cells and outperformed the AAV-2 prototype in tissue in vivo. Vectors with an AAV-DJ capsid were superior in vitro and gave a robust and specific in vivo performance, and provided an ability to evade humoral neutralization by human serum.

IV. Examples

The following examples are illustrative in nature and are in no way intended to be limiting.

Example 1

AAV Capsid Library Generation

A. Plasmids for AAV Capsid Library Generation

Plasmids containing full-length capsid (cap) genes of seven different AAV serotypes were obtained (AAV-2, -4, -5, -8, -9, avian and bovine AAV). Goat AAV was partly synthesized (GeneArt, Regensburg, Germany) as a 888 nt fragment (nt 1023 to 1910). This subclone spans the entire right half of the goat AAV capsid protein, which comprises all 42 reported differences between goat AAV and AAV-5. The other seven cap genes were initially amplified via PCR and subcloned into pBlueScript II SK (Stratagene). The purpose was to flank all cap genes with sites for the unique restriction enzymes Pac I (5') or Asc I (3'), to facilitate later cloning of "shuffled" cap genes into a wildtype AAV plasmid (see below). All primers also contained either a Hind III (5') or a Spe I (3') site, to allow directed cloning into pBlueScript (none of the four restriction enzymes cuts in any parental cap gene). A 20 nt signature region was inserted between the two restriction sites in each primer, to provide conserved primer binding sites for later PCR amplification of shuffled genes. Together, the sequence of the forward primers was 5' GGACTC AAGCTT GTCTGAGTGACTAGCATTCG TTAATTAA CAGGT ATG 3' (SEQ ID NO:6; Hind III site in bold, Pac I site in italics/bold, signature region underlined) directly attached at the 3' end to the first 22 nt of each cap gene following its ATG start codon. Likewise, the reverse primer was 5' CGTGAG ACTAGT GCTTACTGAAGCTCACTGAG GGCGCGCC TTA 3' (SEQ ID NO:7; Spe I site in bold, Acs I site in italics/bold, signature region underlined) directly attached at the 3' end to the last 22 nt of each cap gene up to the TAA stop codon.

In parallel, a wildtype cap recipient plasmid was engineered to contain the AAV-2 packaging elements (ITRs) flanking the AAV-2 rep gene (encoding AAV replication proteins), together with Pac I and Asc I sites for cap cloning, and the AAV-2 polyadenylation site. Therefore, AAV-2 rep (nt 191 to 2189) was PCR amplified using primers containing Bgl II sites and then subcloned into pTRUF3 (carrying AAV-2 ITRs with adjacent Bgl II sites). The forward primer used was 5' CGAACC AGATCT GTCCTGTATTAGAGGTCACGTGAG 3' (SEQ ID NO:8; Bgl II site in bold, AAV-2 nt 191 underlined), and the reverse primer was 5' GGTAGC AGATCT GTTCGACCGCAGCCTTTTGAATGTCCGG TTTATT GATTA GGCGCGCC CTGGACTC TTAATTAA CATTATTGTTCAAAGATGC 3' (SEQ ID NO:9; Bgl II site in bold, polyadenylation signal underlined, Asc I site in italics/bold, Pac I site in italics/bold/underlined, AAV-2 rep stop codon

151737PRTArtificial SequenceSynthetic capsid protein 1Met Ala Ala Asp Gly Tyr Leu Pro Asp
 Trp Leu Glu Asp Thr Leu Ser 1 5 10 15Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro
 Pro Pro 20 25 30Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro 35 40 45Gly
 Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60Val Asn Glu Ala Asp Ala
 Ala Ala Leu Glu His Asp Lys Ala Tyr Asp65 70 75 80Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr
 Leu Lys Tyr Asn His Ala 85 90 95Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly
 Gly 100 105 110Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro 115 120
 125Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140Pro Val Glu His
 Ser Pro Val Glu Pro Asp Ser Ser Ser Gly Thr Gly145 150 155 160Lys Ala Gly Gln Gln Pro Ala Arg
 Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175Gly Asp Ala Asp Ser Val Pro Asp Pro Gln Pro Ile
 Gly Glu Pro Pro 180 185 190Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ala Gly Gly Gly
 195 200 205Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser 210 215 220Ser
 Gly Asn Trp His Cys Asp Ser Thr Trp Met Gly Asp Arg Val Ile225 230 235 240Thr Thr Ser Thr
 Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255Tyr Lys Gln Ile Ser Asn Ser Thr Ser
 Gly Gly Ser Ser Asn Asp Asn 260 265 270Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp
 Phe Asn Arg 275 280 285Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn 290
 295 300Asn Trp Gly Phe Arg Pro Lys Arg Leu Ser Phe Lys Leu Phe Asn Ile305 310 315 320Gln
 Val Lys Glu Val Thr Gln Asn Glu Gly Thr Lys Thr Ile Ala Asn 325 330 335Asn Leu Thr Ser Thr Ile
 Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu 340 345 350Pro Tyr Val Leu Gly Ser Ala His Gln Gly
 Cys Leu Pro Pro Phe Pro 355 360 365Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu
 Asn Asn 370 375 380Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe385 390
 395 400Pro Ser Gln Met Leu Lys Thr Gly Asn Asn Phe Gln Phe Thr Tyr Thr 405 410 415Phe Glu
 Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu 420 425 430Asp Arg Leu Met Asn Pro
 Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser 435 440 445Arg Thr Gln Thr Thr Gly Gly Thr Thr Asn Thr
 Gln Thr Leu Gly Phe 450 455 460Ser Gln Gly Gly Pro Asn Thr Met Ala Asn Gln Ala Lys Asn Trp
 Leu465 470 475 480Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Ser Ala Asp 485 490
 495Asn Asn Asn Ser Glu Tyr Ser Trp Thr Gly Ala Thr Lys Tyr His Leu 500 505 510Asn Gly Arg
 Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His 515 520 525Lys Asp Asp Glu Glu Lys Phe
 Phe Pro Gln Ser Gly Val Leu Ile Phe 530 535 540Gly Lys Gln Gly Ser Glu Lys Thr Asn Val Asp Ile
 Glu Lys Val Met545 550 555 560Ile Thr Asp Glu Glu Glu Ile Arg Thr Thr Asn Pro Val Ala Thr Glu
 565 570 575Gln Tyr Gly Ser Val Ser Thr Asn Leu Gln Arg Gly Asn Arg Gln Ala 580 585 590Ala
 Thr Ala Asp Val Asn Thr Gln Gly Val Leu Pro Gly Met Val Trp 595 600 605Gln Asp Arg Asp Val
 Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro 610 615 620His Thr Asp Gly His Phe His Pro Ser Pro
 Leu Met Gly Gly Phe Gly625 630 635 640Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro
 Val Pro 645 650 655Ala Asp Pro Pro Thr Thr Phe Asn Gln Ser Lys Leu Asn Ser Phe Ile 660 665
 670Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu 675 680 685Gln Lys Glu Asn
 Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser 690 695 700Asn Tyr Tyr Lys Ser Thr Ser Val Asp
 Phe Ala Val Asn Thr Glu Gly705 710 715 720Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu
 Thr Arg Asn 725 730 735Leu22215DNAArtificial SequenceSynthetic capsid protein encoding
 sequence 2atggctgccg atggttatct tccagattgg ctcgaggaca ctctctctga aggaataaga 60cagtgggtgga
 agctcaaacc tggcccacca ccaccaaagc cgcagagcg gcataaggac 120gacagcaggg gtctgtgtct
 tcttggttac aagtacctcg gaccctcaa cggactcgac 180aagggagagc cgtcaacga ggcagacgcc
 gcggcctcg agcacgaca agctacgac 240cggcagctcg acagcgaga caaccgtac ctaagtaca
 accacgccga cgccgagttc 300caggagcggc tcaaagaaga tacgtctttt gggggcaacc tcgggcgagc
 agtctccag 360gcaaaaaga ggcttctga acctcttgt ctggttgagg aagcggctaa gacggctct
 420ggaagaaga ggctgtaga gcactctct gtggagccag actctctc ggaaccgga 480aaggcgggcc
 agcagcctgc aagaaaaga ttgaatttg gtcagactgg agacgcagac 540tcagtccag accctcaacc
 aatcggagaa cctcccag cccctcagg tgtgggatct 600cttacaatgg ctgcaggcgg tggcgacca

